arXiv:1702.07882v1 [math.AT] 25 Feb 2017

DIJKGRAAF-WITTEN Z_2-INVARIANTS FOR SEIFERT MANIFOLDS

SIMON KING, SERGEI MATVEEV, VLADIMIR TARKAEV, AND VLADIMIR TURAEV

Abstract. In this short paper we compute the values of Dijkgraaf-Witten invariants over Z_2 for all orientable Seifert manifolds with orientable bases.
Mathematics Subject Classification 2000: 57M25, 57M27

Seifert manifold, Dijkgraaf-Witten invariant

1. The Dijkgraaf-Witten invariants

In 1990, Dijkgraaf and Witten [DW] proposed a new approach to constructing invariants of closed topological manifolds. Each DW-invariant of a closed oriented n-dimensional manifold M is determined by a choice of a finite group G, a subgroup U of the unitary group $U[1]$, and an element h of the cohomology group $H^n(B;U)$, where $B = B(G)$ is the classifying space of G. Let $S = S(M,B)$ be the set of all base point preserving maps $M \to B$ considered up to base point preserving homotopy. The set S is finite and can be identified with the set of homomorphisms $\pi_1(M) \to G$.

Definition 1.1. The Dijkgraaf-Witten invariant of M associated with h *is the complex number*

$$Z(M,h) = \tfrac{1}{|G|} \sum_{f \in S(M,B)} \langle f^*(h), [M] \rangle,$$

where $|G|$ *is the order of* G *and* $[M]$ *is the fundamental class of* M*.*

In this paper, we consider only the special case where n = 3, both groups G and U have order 2 and are identified with the group $\mathbb{Z}_2$. For the classifying space of $\mathbb{Z}_2$ we take the infinite-dimensional projective space RP^∞, and for h we take a unique nontrivial element $\alpha^3 \in H^3(RP^\infty;\mathbb{Z}_2)$, where α is the generator of $H^1(RP^\infty;\mathbb{Z}_2)$. In this situation, the value $\langle f^*(h), [M] \rangle$ belongs to $\mathbb{Z}_2$, and the formula given above takes the form

$$Z(M,\alpha^3) = \tfrac{1}{2} \sum_{f \in S(M,B)} (-1)^{\langle f^*(\alpha^3), [M] \rangle},$$

and becomes applicable to nonorientable manifolds. If $H^1(M;\mathbb{Z}_2) = 0$, then $Z(M,\alpha^3) = \frac{1}{2}$. In all other cases $Z(M,\alpha^3)$ is an integer.

2. Quadratic function and Arf-invariant

Let M be a closed 3-manifold. We define a quadratic function $Q_M \colon H^1(M;\mathbb{Z}_2) \to \mathbb{Z}_2$ by the rule $Q_M(x) = \langle x^3, [M] \rangle$, where $x \in H^1(M;\mathbb{Z}_2)$, $[M]$ is the fundamental class of M, and $x^3 \in H^3(M;\mathbb{Z}_2)$ is the cube of x in the sense of multiplication in cohomology. The corresponding pairing $\ell_M \colon H^1(M;\mathbb{Z}_2) \times H^1(M;\mathbb{Z}_2) \to \mathbb{Z}_2$ defined by $\ell_M(x,y) = Q_M(x+y) - Q_M(x) - Q_M(y)$ is bilinear.

The following relation between the DW-invariant of M and the Arf-invariant of Q_M was discovered in [MT].

Theorem 2.1. [MT]. *Let M be a closed connected 3-manifold, and let $A \subset H^1(M;\mathbb{Z}_2)$ be the annihilator of ℓ_M. If there exists $x \in A$ such that $x^3 \neq 0$, then $Z(M,\alpha^3) = 0$. If there are no such elements, then $Z(M,\alpha^3) = 2^{k+m-1}(-1)^{\mathrm{Arf}(Q_M)}$, where m is the dimension of A and k equals half the dimension of the coset space $H^1(M;\mathbb{Z}_2)/A$.*

Note that this theorem is true for orientable and non-orientable 3-manifolds. However, it follows from the Postnikov Theorem [Po] that if M is orientable, then the annihilator A of ℓ_M coincides with $H^1(M;\mathbb{Z}_2)$. Further on we will consider only orientable 3-manifolds. For brevity we will call an element $x \in H^1(M;\mathbb{Z}_2)$ *essential* if $x^3 \neq 0$.

Corollary 2.2. *Let M be an orientable closed connected 3-manifold. If there exists an essential $x \in H^1(M;\mathbb{Z}_2)$, then $Z(M,\alpha^3) = 0$. If there are no such elements, then $Z(M,\alpha^3) = 2^{m-1}$, where m is the dimension of $H^1(M;\mathbb{Z}_2)$.*

In view of the above corollary, the following question is crucial for computing DW-invariants: given a 3-manifold M, does $H^1(M;\mathbb{Z}_2)$ contain an essential element? In general, the calculation of products in cohomology and calculation of DW-invariants is quite cumbersome, see [Wa, Br, Ha]. We prefer a very elementary method based on a nice structure of skeletons of Seifert manifolds. For simplicity, we restrict ourselves to Seifert manifolds fibered over S^2, although Theorems 2 – 4 remain true for Seifert manifolds fibered over any closed orientable surface. Proofs are the same.

3. Skeletons of Seifert manifolds.

Definition 3.1. *Let M be a closed 3-manifold. A 2-dimensional polyhedron $P \subset M$ is called* a skeleton of M *if $M \setminus P$ consists of open 3-balls.*

Let us construct a skeleton of an orientable Seifert manifold $M = (S^2; (p_i, q_i), 1 \leq i \leq n)$ fibered over S^2 with exceptional fibers of types (p_i, q_i). Represent S^2 as a union of two discs Δ, D with common boundary. Choose inside Δ disjoint discs $\delta_i, 1 \leq i \leq n$, and remove their interiors. The resulting punctured disc we denote Δ_0. Then we join the circles $\partial\delta_i$ with ∂D by disjoint arcs $l_i \subset \Delta_0, 1 \leq i \leq n$. The skeleton $P \subset M$ we are looking for is the union of the following surfaces, see Fig.1.

(1) Annuli $L_i = l_i \times S^1$ and tori $t_i = \partial\delta_i \times S^1, 1 \leq i \leq n$.
(2) The torus $T = \partial D \times S^1$, the punctured disc Δ_0, and the disc D.
(3) Discs d_i attached to t_i along simple closed curves in t_i of types (p_i, q_i).

Note that the surfaces of the first two types lie in $S^2 \times S^1$ while d_i do not.

4. Seifert manifolds having $Z(M,\alpha^3) = 0$

In this section we describe all orientable Seifert manifolds fibered over S^2 whose first cohomology group contains an essential x. Note that by Poincaré duality $x \in H^1(M;\mathbb{Z}_2)$ is essential if and only if its dual $x_* \in H_2(M;\mathbb{Z}_2)$ is essential in the sense that it can be realized by an odd surface, i.e. by a closed surface having an odd Euler characteristic. So instead of looking for essential 1-cocycles we will construct essential 2-cycles.

Let $M = (S^2; (p_i, q_i), 1 \leq i \leq n)$ be an orientable Seifert manifold fibered over S^2 with exceptional fibers of types (p_i, q_i).



FIGURE 1. A skeleton of M.

Theorem 4.1. *Suppose that M contains an exceptional fiber f_i of type (p_i, q_i) and an exceptional fiber f_j of type (p_j, q_j) such that p_i is divisible by 4 while p_j is even but not divisible by 4. Then there is an essential $x \in H^1(M; \mathbb{Z}_2)$.*

Proof. Choose disjoint discs $D_i, D_j \subset S^2$ containing projection points of f_i, f_j, and join their boundaries by a simple arc $c \subset S^2$. Let P be the polyhedron in M consisting of the annulus $L = c \times S^1$, two tori $t_i = \partial D_i \times S^1, t_j = \partial D_j \times S^1$, and meridional discs $d_i, d_j \subset M$ of solid tori which replace $D_i \times S^1$ and $D_j \times S^1$ by the standard construction of M. The boundary curves of d_i, d_j are of types (p_i, q_i), (p_j, q_j), respectively. See Fig. 2, to the left.



FIGURE 2. Construction of an odd surface.

Note that d_i can be chosen so that the circles ∂d_i and $\lambda_i = \partial L \cap t_i$ decompose t_i into p_i quadrilaterals admitting a black-white chessboard coloring. The same is true for d_j and t_j. Let us remove from P all $(p_i + p_j)/2$ quadrilaterals having the same (say, white) color. We get a closed surface $F \subset M$. Since the Euler

characteristic $\chi(P)$ of P is even and $(p_i + p_j)/2$ is odd, $\chi(F)$ is odd. Therefore F is odd and thus represents an essential 2-cycle. □

Theorem 4.2. *Let $M = (S^2; (p_i, q_i), 1 \le i \le n)$ be an orientable Seifert manifold fibered over S^2 with exceptional fibers of types (p_i, q_i) such that (1) all p_i are odd and (2) the number of odd q_i is even. Denote by Q_* the sum of all q_i, and by P_* any alternating sum of all p_i for which the corresponding q_i are odd. Suppose that the integer number $\xi(M) = (Q_* + P_*)/2$ is odd. Then there exists an essential $x \in H^1(M; \mathbb{Z}_2)$.*

Proof. Let us replace two exceptional fibers of types (p_i, q_i), (p_j, q_j) by exceptional fibers of types $(p_i, q_i + p_i)$, $(p_j, q_j - p_j)$. These operations (called *parameter trading*) preserve the parity of $\xi(M)$ and produce another Seifert presentation of M, which also satisfies assumptions (1) and (2) of Theorem 4.2. Using such operations one can easily get all q_i be divisible by 4, except the last one, say, q_n, which must be even in view of assumption (2). For this new presentation of M we have $Q_* \equiv q_n \mod 4$ (since all other q_i are divisible by 4), and $P_* = 0$ (since there are no odd q_i). Taking into account that $\xi(M)$ is odd, we may conclude that $q_n/2$ is odd. Let P be a skeleton of M constructed for this new presentation of M. Consider the union P_0 of the following 2-components of P, see Fig. 2, to the right:

(1) The torus t_n and disc d_n attached to t_n along a simple closed curve of type (p_n, q_n);
(2) The disc $D = S^2 \setminus \mathrm{Int}(\delta_n)$, where $\delta_n \subset S^2$ is the meridional disc of t_n.

Then we apply the same trick as in the proof of Theorem 4.1 using the circle ∂d_n and taking the circle $\partial \delta_n$ instead of λ_n.

Since q_n is even, the circles ∂d_n and $\partial \delta_n$ can be chosen so that they decompose t_n into q_n quadrilaterals admitting a black-white chessboard coloring. Let us remove from P_0 all $q_n/2$ quadrilaterals having the same (say, white) color. We get a closed surface $F \subset M$. Since the Euler characteristic of P_0 is even and $q_n/2 = \xi(M)$ is odd, $\chi(F)$ is odd. Therefore F represents an essential 2-cycle. □

Let us introduce two classes $\mathcal{A}, \mathcal{B}$ of Seifert manifolds. Class $\mathcal{A}$ consists of manifolds satisfying assumptions of Theorem 4.1, class $\mathcal{B}$ consists of manifolds satisfying assumptions of Theorem 4.2.

Theorem 4.3. *Let M be an orientable Seifert manifold fibered over S^2 with exceptional fibers of types $(p_i, q_i), 1 \le i \le n$. Suppose that M contains an essential 2-cycle x_*. Then M belongs either to $\mathcal{A}$ or to $\mathcal{B}$*

In the proof of the theorem, we need the following self-evident lemma.

Lemma 4.4. *Let τ be a torus and $\mathcal{C}$ a finite collection of simple closed curves in τ such that all their intersection points are transverse. We will consider the union G of these curves as a graph. Suppose that the faces of G have a black-white chess coloring in the sense that any edge of G separates a black face from a white one. Then any general position simple closed curve in τ crosses the edges of G at an even number of points.* □

Proof of Theorem 4.3. Let P be the skeleton of M constructed in section 3. Denote by Γ its singular graph consisting of triple and fourfold lines of P. The remaining part of P consists of 2-components of P, that is, of surfaces which are glued to

Γ along their boundary circles. Let B be the carrier of x_*, i.e. the union of 2-components of P which are black in the sense that they have coefficient 1 in the 2-chain representing x_*. All other 2-components are white.

Case 1. Suppose that Δ_0 and hence D are white. Then the polyhedron $P_0 = P \setminus \Delta_0$ is simple in the sense that the set of its singular points consists of triple lines and their crossing points. It follows that B is a closed surface. Since x_* is essential, $\chi(B)$ is odd. We may assume that B is connected (if not then the Euler characteristic of at least one component of B is odd, and we can take it instead of B). It follows that B contains two annuli, say, L_i, L_j, discs d_i, d_j, and one of the two annuli into which the curves $L_i \cap T$ and $L_j \cap T$ decompose the torus T. The total Euler characteristic of these 2-components is 0.

Just as in the proof of Theorem 4.1 the circles ∂d_i and $\lambda_i = t_i \cap \partial L_i$ decompose t_i into p_i quadrilateral 2-components. Since x_* is a 2-cycle, they are colored according to the chessboard rule. It follows that p_i and p_j are even. The remaining part of B consists of $(p_i + p_j)/2$ black quadrilaterals in t_i and t_j. Since $\chi(B)$ is odd, we may conclude that one of $p_i/2, p_j/2$ is even while the other is odd. Therefore $M \in \mathcal{A}$.

Case 2. Suppose that Δ_0 is black. Then each d_i is black. Otherwise the boundary of x_* would contain $\partial\delta_i$, which is impossible since x_* is a 2-cycle.

Let us prove that if L_i is black then q_i is odd. Consider the graph $G = \partial d_i \cup \lambda_i \subset t_i$. As above, x_* induces a chess black-white coloring of its faces. Then a parallel copy $\partial\delta_i'$ of $\partial\delta_i$ crosses λ_i at one point and crosses ∂d_i at q_i points. By Lemma 4.4 the number $q_i + 1$ must be even, which means that q_i is odd. Similarly, if L_i is white then q_i is even.

Let us prove that all p_i are odd. Suppose that L_i is black. Consider the graph $G = \partial d_i \cup \partial\delta_i \subset t_i$. It decomposes t_i into black-white colored faces. The coloring is induced by the 2-cycle x_*. Let $\lambda_i' \subset t_i$ be a parallel copy of λ_i. Then λ_i' crosses ∂d_i at p_i points and crosses $\partial\delta_i$ at one point. By Lemma 1 the number $p_i + 1$ must be even, which means that p_i is odd. Suppose that L_i is white then q_i is even. Therefore , p_i, being coprime with Q_i, is odd.

Let us prove that the number of black L_i and hence the number of odd q_i are even. This is because the boundary circles of black L_i decompose T into black-white colored annuli such any two neighboring annuli have different colors. Therefore any meridional circle of T, for example, ∂D, crosses those boundary circles at an even number of points.

Let us prove that $\xi(M)$ is odd. Just as in the proof of Theorem 4.2 we transform the given Seifert presentation of M into a new Seifert presentation such that all $q_i, 1 \le i \le n-1$, are divisible by 4 and q_n is even. Then the carrier B of x_* consists of the following black surfaces:

(1) Δ_0 and D;
(2) All discs $d_i, 1 \le i \le n$;
(3) black quadrilaterals contained in $t_i, 1 \le i \le n$. Each t_i contains $q_i/2$ black quadrilaterals, where (p_i, q_i) is the type of the corresponding d_i. Of course B may contain the torus T, but it is is homologically trivial and thus can be neglected.

Note that B has only triple singularities and thus is a closed surface. Since x_* is essential, $\chi(B)$ is odd. Now we calculate $\chi(B)$ by counting Euler characteristics of the above black surfaces. We get $\chi(B) = q_n/2 \bmod 2$. It follows that $q_n/2$ is odd. Since all q_i are even, $P_* = 0$. Taking into account that all $q_i, 1 \le i \le n-1$

are divisible by 4 and that $q_n/2$ is odd we may conclude that $Q_* = (\sum_{i=1}^n q_i)/2$ is odd. Therefore, $\xi(M)$ is odd. It follows that is in class $\mathcal{B}$. □

Acknowledgments

S. Matveev, V. Tarkaev and V. Turaev were supported in part by the Laboratory of Quantum Topology, Chelyabinsk State University (contract no. 14.Z50.31.0020). S. Matveev and V. Tarkaev were supported in part by the Ministry of Education and Science of the Russia (the state task number 1.1260.2014/K) and the Russian Foundation for Basic Research (project no. 14-01-00441).

References

[Br] Bryden, J., Hayat-Legrand, C., Zieschang, H., Zvengrowski, P. *The cohomology ring of a class of Seifert manifolds, Topology Appl.105 (2) (2000), 123-156.*

[DT] Deloup, F., Turaev, V., *On reciprocity.* J. Pure Appl. Algebra 208 (2007), no. 1, 153–158.

[DW] Dijkgraaf, R., Witten, E., *Topological gauge theories and group cohomology,* Comm. Math. Phys. 129 (1990) 393–429.

[Ha] Haimiao Chen, *The Dijkgraaf-Witten invariants of Seifert 3-manifolds with orientable bases, arXiv:.1307.0364v3 [math.GT] 6Apr 2014.*

[MT] Matveev, S.V., Turaev V.G. *Dijkgraaf-Witten Invariants over Z_2 for 3-Manifolds,* Doklady Mathematics, 2015, Vol. 91,No.1. pp.9-11.

[Po] Postnikov, M. M., *The structure of the ring of intersections of three-dimensional manifolds.* (Russian) Doklady Akad. Nauk. SSSR (N.S.) 61, (1948). 795–797.

[Wa] Wakui, M., *On Dijkgraaf-Witten invariant for 3-manifolds.* Osaka J. Math. 29 (1992), no. 4, 675–696.

Faculty of Mathematics and Natural Science, Institute of Mathematics Education, Gronewaldstr. 2, D-50931 Cologne Germany, simon.king@uni-koeln.de

Laboratory of Quantum Topology, Chelyabinsk State University, Brat'ev Kashirinykh street, 129, Chelyabinsk 454001 Russia, and, Krasovsky Institute of Mathematics and Mechanics of RAS, svmatveev@gmail.com

Laboratory of Quantum Topology, Chelyabinsk State University, Brat'ev Kashirinykh street, 129, Chelyabinsk 454001 Russia, and, Krasovsky Institute of Mathematics and Mechanics of RAS, v.tarkaev@gmail.com

Department of Mathematics Indiana, University Bloomington IN47405, USA, and, Laboratory of Quantum Topology, Chelyabinsk State University,, Brat'ev Kashirinykh street, 129, Chelyabinsk 454001 Russia, vturaev@yahoo.com